



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/82, 15/54, A01H 5/00</b>		<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/09188</b>
			<b>(43) International Publication Date:</b> 25 February 1999 (25.02.99)
<b>(21) International Application Number:</b> PCT/EP98/04988 <b>(22) International Filing Date:</b> 27 July 1998 (27.07.98) <b>(30) Priority Data:</b> 97202507.6 13 August 1997 (13.08.97) EP <b>(71) Applicant (for all designated States except US):</b> VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE [BE/BE]; Rijvisschestraat 118, B-9052 Zwijnaarde (BE). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> CHEN, Cuiying [CN/BE]; Meibloemstraat 143, B-9000 Gent (BE). MEYERMANS, Hugo [BE/BE]; Breugelhoestraat 25, B-2560 Nijlen (BE). VAN MONTAGU, Marc [BE/BE]; De Stassartstraat 120, B-1050 Brussel (BE). BOERJAN, Wout [BE/BE]; Zomerstraat 44 <sup>b</sup> , B-9270 Laarne (BE). <b>(74) Common Representative:</b> VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE; Rijvisschestraat 118, B-9052 Zwijnaarde (BE).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> TISSUE-SPECIFIC POPLAR PROMOTERS			
<b>(57) Abstract</b>  The invention concerns the isolation and characterization of DNA sequences representing a caffeoyl-CoA-O-methyltransferase (CCoAOMT) promoter having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel or vessels. The cells adjacent to the vessel are xylem ray cells whereas the vessel can be a differentiating vessel.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## Tissue-specific poplar promoters

The present invention relates to isolated DNA sequences representing a *CCoAOMT* promoter or a functional part thereof having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel.

### Background description to the invention

After cellulose, lignin is the major structural component of secondary thickened plant cell walls. It plays a very important role in vascular plants; it provides rigidity to the cell wall and hydrophobicity to the water-conducting vascular elements. Moreover lignin is often deposited after pathogen attack to prevent the spread of a pathogen. Lignin is a complex polymer of hydroxylated and methoxylated monolignol units that are termed *p*-hydroxyphenyl (H), guaiacyl (G), and syringyl (S). These units differ in their degree of methoxylation: the H unit is not methoxylated at the C5 position, the G unit once, and the S unit twice (see Figure 1, lignin biosynthesis pathway). These units are subsequently dehydrogenated by peroxidase(s) and possibly laccase(s) to form mesomeric radicals which polymerise to form the complex lignin network (Boudet et al., 1995). Gymnosperm lignin consists mainly of G units. Angiosperm dicot lignin incorporates both G and S units whereas grass lignin consists of all three units. The S/G ratio impacts the physical properties of lignin. Lignin content increases with maturity in stems (Jung and Vogel, 1986), and the composition changes with advanced maturity towards a progressively higher S/G ratio (Buxton and Russell, 1988). In the paper industry, it has been long a problem to remove lignin from cellulose by chemical paper pulping, because of the huge cost and the large environmental pollution (Tien, 1987). Both a high lignin concentration (Albrecht et al., 1987) and a low S/G ratio have a negative impact on lignin extractability during chemical pulping. Lignin also limits forage digestibility. Even a small decrease in lignin content has a significant positive impact on paper pulping and digestibility and

therefore profitability (Casler, 1987). Therefore, there is considerable interest in the potential for genetic manipulation of the lignin level and/ or composition to improve digestibility of forages and pulping properties of trees (Dixon et al., 1994).

The biosynthesis pathway of lignin precursors proceeds through the common phenylpropanoid pathway starting from phenylalanine and leading to the synthesis of cinnamoyl-CoAs. Subsequently, the cinnamoyl-CoA esters are channelled into the lignin branch pathway to produce cinnamyl alcohols. The methylation of the 3- and / or 5-hydroxyl group of hydroxycinnamic acids is an important step influencing lignin composition because the chemical structures of the monomeric lignin precursors differ only in 3- and / or 5-methoxyl groups on the aromatic ring. The O-methyltransferases (OMTs) involved in lignin formation have been characterised in a number of species (Grisebach, 1981; Bugos et al., 1991; Gowri et al., 1991; Collazo et al., 1992; Davin and Lewis, 1992). Caffeic acid O-methyltransferase (COMT, EC2.1.1.68) has long been considered as the sole methylation pathway involved in lignification (Neish, 1968; Grisebach, 1981; Lewis and Yamamoto, 1990). However, there is now increasing evidence suggesting that the O-methylation of the lignin precursors may also occur at the level of the hydroxycinnamoyl-CoA esters. A specific O-methyltransferase involved in the methylation of caffeoyl-CoA (caffeoyl-Coenzyme A 3-O-methyltransferase; CCoAOMT) was initially characterised in parsley cell suspensions (Kneusel et al., 1989) and in carrot cell suspensions (Kuhn et al., 1989). In parsley cell suspension cultures, the addition of a fungal elicitor induced both *p*-coumaroyl-CoA 3-hydroxylase (CCoA-3H) and CCoAOMT activities. The resulting feruloyl-CoA however, first was thought to be incorporated uniquely into cell wall-bound feruloylated polymers implicated in the defence response. However, Ye et al (1994) and Ye and Varner (1995) recently suggested that the CCoAOMT mediated pathway is dominant in lignifying *Zinnia* tracheary elements (TES) *in vitro*; they showed that CCoAOMT, but not COMT is induced during tracheary element formation. Ye and Varner (1995) suggested that, like in White birch and *Arabidopsis*, tracheary elements may be composed of mainly G units and that therefore COMT might not be expressed under these conditions. Nevertheless, although in these published works CCoAOMT was suggested to be involved in lignification, a direct link has still to be found. Furthermore, it was found that in COMT down-regulated tobacco (Atanassova et al, 1995) and poplar (Van Doorselaere et al, 1995) the S/G ratio was

decreased. The modification of the lignin composition in these transgenic plants was mainly due to a decrease in the number of S units and an increase in the level of G units. These results demonstrate that the activity of COMT mainly controls the level of S units and therefore it was hypothesised that the production of G units in COMT down-regulated plants may result from the functioning of a CCoAOMT-mediated methylation pathway.

#### Detailed description of the invention.

The present invention concerns the isolation and characterisation of two genes encoding CCoAOMT in poplar which share high homology in the coding region but are diverse in their regulatory parts. To study its specific expression at the cellular level in plants, histochemical analysis have been performed in transgenic poplars harbouring chimeric pCCoAOMT-GUS constructs. In addition, the CCoAOMT protein has been immunolocalised. The data show that both genes are differentially expressed in phloem fibers. In the xylem of the stem, both genes are expressed preferentially in the ray cells next to vessel elements for which it is known that the lignin is mainly composed of G units. These data show that CCoAOMT is involved in lignification for providing G units to the cell wall, even after autolysis of the cytoplasm. The invention further relates to an isolated DNA sequence according to figure 1 B (also called SEQ ID NO 1) which represents a CCoAOMT promoter or a functional part thereof having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel.

Furthermore an isolated DNA sequence according to figure 1 C (also called SEQ ID NO 2) which represents a CCoAOMT promoter or a functional part thereof having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel belongs to the current invention.

The cells adjacent to the vessel as defined can be xylem ray cells and xylem fibers whereas in addition the vessel can be a differentiating vessel.

Part of the invention is a promoter sequence conferring expression in living vessel elements and/or in ray cells, adjacent to living or autolysed vessel elements and/or in fibers adjacent to said living or autolysed vessel elements, with the proviso that said

expression does not occur constitutively in xylem fiber cells or ray cells that are not adjacent to said vessel elements.

In this last described situation the expression conferred by the promoter or a functional part thereof according to the invention occurs preferably in those ray cells comprising a so-called pith to the vessel as is clearly demonstrated in figures 5A and 5B respectively.

With "a functional part of a promoter" is meant in this description, any part or a fragment of a promoter that can still induce gene expression in the cells as disclosed herein. Such a fragment or part can be used alone or in combination with a second promoter and/or parts or fragments of said second promoter.

With "gene expression" is meant a sequence of events that results in the synthesis of RNA starting from DNA (so called transcription) regardless whether or not the resulting RNA is translated into a protein.

Furthermore, the expression of both above mentioned chimeric genes was markedly induced by wounding and fungal infection. GUS analysis showed that the expression of both chimeric genes was only induced when lignin was deposited at the wounded zone. The similar correlation between the induced expression of chimeric genes and the induced formation of lignin was detected at the site of fungal infection as well. So it has been shown that the regulation of both chimeric *CCoAOMT* genes is involved in lignification not only during normal development but also in response to wounding and pathogen infection.

To the scope of the invention also belongs a recombinant DNA comprising any of the isolated DNA sequences according to the invention whereas an embodiment of the current invention is represented by a plant or tree cell comprising said recombinant DNA integrated in its genome. Furthermore to the invention belongs a transgenic plant or tree comprising said cell and its progeny thereof but also seed, seedlings, roots and the like.

In order to localize promoter regions potentially involved in the transcriptional control of the *CCoAOMT* gene, several deletions were generated by removing part of the 5' flanking sequences of both *CCoAOMT* genes (Fig. 14), and examined their effect on gene expression in transgenic poplars. 4-13 individual transformants per construct were analyzed.

For the deletions of the *PtCCoAOMT1* gene, differences were found in the tissue specific pattern of GUS expression between plants containing the full-length promoter (*pBINPOP1*) and the various promoter deletions. As indicated in Table 3 GUS activity was undetectable in poplar transformed with the *pBIN1DD1* (-114) promoter-GUS construct. In poplar transformant with the deletion *pBIN1DC2* (-184) no GUS activity could be detected in the xylem tissue, whereas faint GUS activity was detected in phloem fibres, cortex and periderm (Fig. 15). The fact that the *pBIN1DC2* (-184) is sufficient for directing expression in bark tissue, a *cis*-positive regulatory element (domain +B) could lie between -184 and -114 bp upstream of the translation start site (Fig. 17). Construct of *pBIN1DB3* (-199) which is 15 bp longer than *pBIN1DC2* failed to direct expression in any tissue, suggesting that this additional 15 nucleotides (which includes an AC-II element) contains a *cis*-negative regulatory element (domain -B) for controlling the expression in bark tissue. In the transformants containing *pBIN1DA5* (-456), GUS staining remained in the xylem vessels and xylem ray cells adjacent to these vessels and in the periderm as for the full-length promoter *pBINPOP1*. However, it was found that *pBIN1DA5* (-456) lost expression in companion cells and conferred additional expression in cambial ray initial cells (Fig.16). This observation suggested the presence of two domains between -456 and -199 bp upstream of the translation start site of the *PtCCoAOMT1* promoter: one domain (+CR) positively controls expression in cambial ray cells, the other domain (V) positively regulates expression in vessels and adjacent ray cells. Furthermore, the comparison between the expression patterns conferred by the full-length promoter *pBINPOP1* and *pBIN1DA5* suggests the presence of a domain (CC) that directs expression in companion cells and a domain (-CR) that negatively controls the expression of *CCoAOMT* in cambial ray cells between -1993 and -456 bp upstream of the translation start site of the *PtCCoAOMT1* promoter.

For the *PtCCoAOMT2* promoter, the deletion of *pBIN2DD1* (-110) and *pBIN2DB3* (-195) resulted in a complete loss of expression. *PBIN2DA4* (-497) however directed expression as the full-length *PtCCoAOMT2* promoter. The -497 fragment of *PtCCoAOMT2* promoter is sufficient to control the cell-specific expression pattern, suggests that *cis*-acting positive regulatory elements, which control the expression in xylem vessels and in adjacent ray cells (domain V) and in phloem fibres (domain F), are located between nucleotides -497 and -195.

The transgenic poplars containing the promoter deletions were also analyzed upon wounding. The GUS staining revealed that *pBIN1DA5* and *pBIN2DA4* were significantly expressed at the wounding zone of leaves 3-5 days post wounding as their full-length promoters. The P-HCl staining indicated that lignin was deposited at the site of wounding as well. The other deletions did not show any GUS activity at the wounding zone, suggesting that a *cis*-acting positive regulatory element (domain W) for response to wounding is located between -456 and -199 of the *PtCCoAOMT1* promoter and between -497 and -195 of the *PtCCoAOMT2* promoter.

Taken together, the two full-length *CCoAOMT* promoters and their deletions revealed several domains of importance for regulated expression, as summarized in Table 3. The putative regions that could contain *cis*-acting element(s) for cell- and tissue-specific expression are summarized in Figure 17. By nucleotide sequence comparison of *CCoAOMT* promoters from different species several conserved regions were identified between the different sequences of each promoter and between the both promoters (Fig. 13). These regions are probably *cis*-regulatory elements for controlling the cell- and tissue-specific expression of *CCoAOMT*.

The invention is further explained by non-limiting examples given hereunder.

## EXAMPLES

### **1. Genomic clones and copy number of *CCoAOMT* genes**

Genomic clones were isolated from a poplar genomic library (*P. trichocarpa* cv Trichobel), by using a cDNA encoding *CCoAOMT* from poplar (*P. trichocarpa* cv Trichobel) as a probe. Six individual positive plaques were screened, and two of them, which are designated as *Pop1* and *Pop2*, respectively, were studied in detail (Fig. 1 A-D). Comparison of the nucleotide sequences of these two genes, 3.8kb and 3kb in length respectively, to the *CCoAOMT* cDNA shows that both genes consist of five exons and four introns with identical intron locations. Both deduced polypeptides have 340 aa, with a calculated MW of 27.8 kDa for *Pop1*, and 27.9 kDa for *Pop2*. Alignment of the two genes revealed that they share high identity in the coding regions (93.9% at amino acid level), but are significantly divergent in the noncoding parts. In *Pop1*, a putative polyadenylation signal AATAAA is found 82 nucleotides



downstream from the TGA stop codon whereas in *Pop2*, an ATAGTT sequence is located at 68 nucleotides downstream of the stop codon. The transcription start site was determined by comparison with the corresponding cDNA sequences. The TATA motif of *Pop1* and *Pop2* was found at position 125 bp and 121 bp upstream from the translation start site ATG, respectively. According to the *CCoAOMT* cDNA sequence, the transcription start site is situated at 71 bp upstream the translational start point in *Pop1* and 69 bp in *Pop2*. Several conserved regions were found in the promoters of these two genes.

## 2. Expression patterns conferred by the *CCoAOMT* promoters

In order to study the tissue and cell specific expression of both genes, the two promoters of *Pop1* and *Pop2*, 2kb and 1.3 kb in length, respectively, were fused to the  $\beta$ -glucuronidase (GUS) reporter gene (Figure 2). Poplar (*P. tremula* x *P. alba*) was stably transformed with both chimeric constructs via *Agrobacterium tumefaciens*-mediated transformation. Histochemical GUS assays were done on three-month-old greenhouse-grown transgenic poplars.

Figure 3 shows a transversal section through a young internode, stained with phloroglucinol-HCl for lignin, and analysed for GUS activity. The primary xylem stains red. At this stage GUS activity, driven by the two promoters according to the invention, was detected in the xylem ray cells between the vessels, in the pith parenchyma cells surrounding the inner primary xylem vessels, and within developing vessels themselves (Fig. 4). No GUS staining can be seen in the lignified parts of primary xylem, between the vessel bundles (Fig. 3 and 4).

In older internodes, GUS activity conferred by the two promoters is present in the secondary xylem shown in a cross section through the stem (Fig. 5). Observations under higher magnification of the xylem tissue clearly demonstrated that GUS activity was preferentially localised in xylem ray cells just next to vessels and within young developing vessels themselves (Fig. 5 and 6). Most expression was detected in the younger parts of the xylem, less towards the inner part of the xylem. No GUS activity was detected in xylem fibers throughout the whole stem, except for fibers that were adjacent to vessels (see further).

Although both promoters gave a very similar expression pattern in the xylem, they were differentially regulated in phloem tissue. *Pop1* conferred expression

specifically in companion cells of the phloem and in the cells surrounding the phloem fibers, whereas *Pop2* conferred expression within the phloem fiber cells (Fig. 7 and 8).

A similar cell specific expression pattern was detected in petioles and leaves (Fig. 9 and 10); GUS analyses revealed that both chimeric genes had similar expression patterns in the xylem of these organs, but had a different expression pattern in the phloem. *Pop2* conferred expression preferentially in phloem fibers (Fig. 9B and 10B) while *Pop1* preferentially in phloem parenchyma cells (Fig. 9A and 10A).

Figures 11A and 11B show that both chimeric genes were similarly expressed in xylem ray cells next to vessels and in phloem fibers in roots.

In addition, both chimeric genes were strongly expressed in the phellogen and phelloderm of the periderm. The blue staining of the GUS activity was localised in the cell layer adjacent to the cell layer that stained red with phloroglucinol-HCl (Figure 7 and 8). GUS activity was also detected in the meristem (figure 11C).

### 3. Response to mechanical wounding

The biosynthesis of lignin is not only an essential process under normal conditions but is also believed to be crucial in responses to stress such as mechanical wounding (Matern and Grimmig, 1994; Nicholson and Hammerschmidt, 1992).

To study the involvement of *CCoAOMT* in response to mechanical wounding during the formation of a lignin barrier, the spatio-temporal expression of the two chimeric genes was histochemically characterised in leaf tissue at different periods after wounding. Histochemical assays in combination with GUS staining and phloroglucinol-HCl staining revealed that no GUS activity nor lignified cells were detected in leaf samples harvested one, two, three and four days after wounding. When the necrophylactic layer became visible surrounding the cutting site in leaf samples five days after wounding, phloroglucinol-HCl staining revealed a barrier, suggesting the deposition of lignin or lignin-like material. At this stage, GUS staining showed that expression conferred by both *Pop1* and *Pop2* promoters was associated with the lignified barrier. When the barrier was well developed in leaf samples harvested seven days after wounding, no GUS activity was detected anymore.

Table 2 summarises the expression characteristics conferred by both promoters.

#### 4. Expression of both genes is induced by fungal infection

To investigate whether expression of *CCoAOMT* is a response to pathogen infection, detached leaves from transgenic and wild type poplar were sprayed with spores of the fungus *Melampsora pinitorca*, a natural pathogen for poplars belongs to *Leuce* section. Three days after infection, *in vitro* leaves, induced GUS activity was observed. Subsequently the infected leaves die within 5 to 10 days without appearing uredosores. In contrast, in the leaves of green house grown plants seven days after infection, the orange uredosores became visible and necrotic lesions developed at the site of infection. At this stage, GUS activity driven by the both promoters was strongly induced in the cells surrounding these uredosores and necrotic lesions, whereas no GUS staining was observed before the formation of uredosers. The blue colour of the staining is closely associated to the lignified region at the infection sites, indicating a correlation between the induced expression of *CCoAOMT* chimeric genes and lignification in necrotic lesions both *in vitro* and *in vivo* leaves.

#### 5. Immunocytochemical localisation of CCoAOMT

In order to confirm the GUS analysis, immunolocalisations were performed using stem sections in which secondary growth was apparent, using polyclonal antibodies for *CCoAOMT*. Immuno-fluorescence microscopy revealed labelled proteins as a fluorescent green colour. *CCoAOMT* was localised intensively within differentiating vessels, and specifically in xylem ray cells just next to lignifying vessels (Figure 12B). In addition, labelling was detected on the secondary thickened walls of the phloem fibers (Figure 12C). No immunolabeling was observed in xylem fibers, except for the fibers that were adjacent to vessel elements. In these fiber cells, immunolabel was detected preferentially at that site of the cell that was adjacent to a vessel. The results indicate that the pattern of localisation of *CCoAOMT* in stem tissue was very similar to the pattern obtained by histochemical GUS assays. Hardly any signal could be observed in sections of the same stem treated with rabbit pre-immune serum (Figure 12A)

#### 6. Cloning and sequencing the promoters from different species of poplar

It was reasoned that the *cis*-elements important for tissue specific expression should be conserved between a series of allelic forms of the promoter itself. In order to identify possible elements responsible for the tissue specific expression pattern, a

PCR-based promoter amplification strategy was developed. By alignment of the two *CCoAOMT* promoters, conserved sequences were identified. These conserved sequences were used to design primers that would allow the amplification of the homologous sequences from other poplar species (Figure 1D). The PCR products were cloned into a pGEM-T vector and sequenced. By comparing these sequences with each other, several conserved sequences, present in all promoters were identified (Fig. 13). The AC-element II, a putative regulatory element involved in UV light and /or elicitor regulation in promoters of other plant phenylpropanoid biosynthesis genes such as PAL and 4CL (Table 1), was conserved in both promoters.

For sake of clarity in order to better understand the invention and the materials used therein a "Material and Method" section is given hereunder.

## **Material and methods**

### **1. Screening of a genomic library**

A lambda FIX II custom genomic library (Stratagene, USA) from poplar (*Populus trichocarpa* cv Trichobel) was screened by using a <sup>32</sup>P-labelled poplar *CCoAOMT* cDNA as a probe. Six positive clones were isolated. Lambda DNA was prepared and digested by various restriction enzymes, and the fragments homologous to the cDNA probe were subcloned into a pBluescript vector (Stratagene).

### **2. DNA sequence analysis**

The nucleotide sequence of genomic subclones in Bluescript and a series of overlapping deletions was determined by dideoxynucleotide sequencing.

### **3. The *Pop-GUS* fusion constructions**

The promoter fragments (*Pop1*, 2kb and *Pop2*, 1.3kb) were translationally fused to the *uid A* coding region in plasmid pGUS1 (Peleman et al., 1989). The fragments containing the chimeric promoter-GUS fusions were cloned into the binary vector pBIN19 (Bevan, 1984).

#### 4. Plant material and transformation

Poplar plants used in stable transformation experiments corresponded to the INRA clone 717-1B4 (*Populus tremula* x *Populus alba*). This line was selected because of its susceptibility to *Agrobacterium tumefaciens* infection (Leplé et al., 1992). The *popGUS* constructs in pBIN19 were mobilised to *Agrobacterium* strain C58C1Rif harbouring the plasmid pMP90 by the freeze-thaw method described by Zham et al. (1984). Poplar was transformed with the chimeric constructs following the approach described by Leplé et al. (1992). *In vitro* plants were maintained on half MS medium at 24 °C with a photo-period of 16 h light and 8 h darkness. About two-month old plantlets were transferred to a greenhouse. Transgenic poplars were grown in the greenhouse at 21 °C with the same light cycle. GUS analyses were performed after three months of growth in the greenhouse.

The following species of poplar were used to make promoter comparisons between different species of poplar: *P.trichcarpa*, *P.euphratica*, *P.laurifolia*, *P. nigra*, *P. deltoides*, *P.alba*, *P. maximowiczii*, and *P. ciliata* (see Figure 13).

#### 5. Histochemical assays

Histochemical staining for GUS activity was performed according to Jefferson et al. (1987). Stems, roots, leaves and petioles from transgenic plants were sectioned with a vibroslicer (Laborimpex, Brussels, Belgium), fixed in 3% glutaraldehyde in 100 mM potassium phosphate buffer, pH 7.0, for 30 min at room temperature. The sections were kept in the same phosphate buffer. GUS staining was carried out by incubating sections with X-Gluc (5-bromo-4-chloro-3-indolyl glucuronide), K<sub>3</sub>Fe(CN)<sub>6</sub>, and K<sub>4</sub>Fe(CN)<sub>6</sub>, for 1-4 h. Staining was allowed to proceed at 37 °C until blue stain developed in the samples (1 to 4 h).

Lignin was revealed by staining with phloroglucinol-HCl according to Speer (1987).

#### 6. Cloning and sequencing of CCoAOMT promoters from different poplar species

Based on a homology sequence alignment between the two CCoAOMT promoters, a 5' primer was chosen from a conserved sequence localised at about 900 bp upstream of the initiation codon and a 3' primer 20 bp downstream of the

initiation codon (Fig.1-D). Genomic DNA, that was prepared according to Shure et al. (1983), was used as template for PCR. PCR reactions were performed using 500 ng of genomic DNA and 10 pmol of each primer. The reaction mixture was first kept at 94 °C for 4 min followed by 30 cycles under following conditions: denaturation at 94 °C for 1 min, annealing at 45 °C for 1 min, extension at 72 °C for 50 seconds on a PCR machine. The PCR product was cloned directly into the pGEM-T vector (Promega).

## 7. Preparation of antibodies directed against CCoAOMT

CCoAOMT was expressed in *E.coli* as a fusion protein using the pGEX (Pharmacia Biotech) expression vector. The fusion protein was purified using the Glutathione Sepharose 4B Redipack columns (Pharmacia Biotech). Purified native fusion protein was used to immunise a rabbit following the standard protocol. After cutting the purified fusion protein with trombine protease, a second rabbit was immunised with a CCoAOMT sample cut out from SDS-PAGE. Western blotting procedures were standard. (Western blots of poplar xylem extracts using the antiserum raised against the fusion protein and against the CCoAOMT gel sample, showed predominantly two bands, +/- 29 kDa and a weaker +/-26 kDa. The upper band corresponded to CCoAOMT as evidenced by microsequencing the corresponding spots on a two-dimensional gel. The nature of the less intense hybridising band of lower molecular weight is still unknown).

## 8. Deletions (5'-upstream) in CCoAOMT promoter-GUS constructs

A series of CCoAOMT promoter fragments were created by PCR using a pair of specific primers which were homologous to either the 5' upstream or the coding region, as shown in Figure 1-B and 1-C. The PCR products were ligated into the pGEM-T vector by using the pGEM®-T Vector System I Kit (Promega) and subsequently translationally fused to the *uidA* gene in plasmid pGUS1 (Peleman et al., 1989) using the restriction sites *NcoI* and *SacI*. The chimeric GUS fusions were subsequently cloned as *XbaI* fragment into pBIN19 (Bevan, 1984; Figure 14). All the plasmids were verified by restriction digestion and sequencing. Those with desired orientation were selected and transferred into *Agrobacterium tumefaciens* by the freeze-thaw method described by Zham et al. (1984).

## 9. Lignin staining

Lignin was revealed by phloroglucinol-HCl (P-HCl) staining according to Speer (1987). Sections and samples were incubated for 2 min in phloroglucinol solution (1% in ethanol/water 92/8 v/v), then mounted in 25% HCl. The Mañle reaction was used to distinguish lignin rich or poor in syringyl monomers (Monties 1987). For the Mañle reaction, samples were immersed for 5 min in 1% KMnO<sub>4</sub>, and rinsed with H<sub>2</sub>O. Subsequently, the samples were destained for 1 min in 25% HCl, washed with H<sub>2</sub>O, and mounted in 32% of NH<sub>4</sub>OH.

## 10. Wounding and pathogen infection

Mechanic wounding was performed on the petioles of three months old greenhouse plant by making half cm long vertical slit with a scalpel. Wounded petiole regions were excised at day 1, 2, 3, 4, 5, and two weeks post wounding. Two months old transgenic plants were grown in Weck-pot *in vitro* condition. A half leaf was cut off using a sterile scalpel, and a 2-mm width leaf tissue from cutting site was cut and stored directly at -70°C as control. The wounded plant was continuously incubated at the same *in vitro* condition. Subsequently, a 2-mm width of wounded leaf tissue from cutting site was collected after wounding for 1, 2, 3, 4, 5 days, respectively. The harvested samples were then used for GUS assays.

For pathogen infection, the detached leaves were sprayed with spores of fungus (*Melampsora pinitorca*) on the under side, in a concentration of 200,000 spores/ml. The infected leaves were floated up side down on water in Petri dishes and were incubated at 22° C in a greenhouse for various periods.

Strictly for the case of clarity some key elements of the current invention are discussed below in more detail.

The tissue and cell specific expression pattern of CCoAOMT has been demonstrated in a woody plant. The GUS reporter system has permitted to reveal the temporal and spatial expression patterns conferred by the *Pop1* and *Pop2* promoters. Histochemical analysis of GUS activity showed that both *Pop1* and *Pop2* were

predominantly expressed in the vascular tissue of stems, leaves, petioles, and roots. The GUS data presented here are consistent with the tissue printing analysis described earlier by Ye and Varner (1995). The chimeric *Pop1-GUS* and *Pop2-GUS* constructs exhibited some different and some overlapping patterns of GUS expression, and these are summarised in Table 2. The tissue-specific expression of the *Pop1* and *Pop2* promoters were strictly correlated with tissues undergoing active lignification such as xylem and phloem fibres.

In the xylem of stems throughout the whole plant, both *Pop1* and *Pop2* promoters conferred expression in the xylem ray cells adjacent to vessels and in differentiating vessel themselves. Immunolocalisation studies confirmed the accumulation of CCoAOMT in the ray cells next to vessels and within differentiating vessels themselves. A strong preferential expression is conferred by the promoters in ray cells adjacent to vessel elements. For many of the lignin biosynthesis genes, the tissue specific expression has been analysed, either by RNA tissue printing, or by promoter-GUS fusions. In bean expression conferred by the *PAL*-promoter was also localised preferentially in xylem ray cells (Bevan et al. 1989). Hauffe et al. (1991) reported the localisation of GUS activity under control of the *4CL* promoter preferentially in xylem at the onset of tracheary element differentiation, and strictly in xylem ray cells positioned between highly lignified tracheary elements (vessels and fibres). Smith et al. (1994) provided further evidence by complementary immunolocalisation, that *PAL* and *4CL* accumulate in cells adjacent to the metaxylem in bean hypocotyls. However, *PAL* and *4CL* are also involved in the synthesis of other phenylpropanoid end products. The role of xylem parenchyma cells in lignin synthesis is still questionable. Recently, Feuillet et al.(1995) and Hawkins (1997) reported that the promoter of *CAD* in *Eucalyptus* was expressed in regions undergoing active lignification i.e. phloem fibres, differentiating xylem, ray parenchyma cells, and vascular cambium. The *CAD* promoter was active throughout the ray, thus next to vessels but also fibers.

It is indicated according to the invention that CCoAOMT promoter activity, unlike those of *PAL*, *4CL* and *CAD* which are constitutively present in xylem ray cells, is strictly located in cells just next to vessels. It is strongly suggested that CCoAOMT is closely correlated to vessel lignification.



It has been shown by Osakabe et al., (1996) that lignin in the cell wall of vessel elements is predominantly G lignin. This is in contrast to the lignin in fibers cell walls, which consists of both G and S units in comparable amounts. Also in white birch and *Arabidopsis*, lignin in vessel cell walls incorporates predominantly G units (Saka and Goring, 1988; Chapple et al., 1992). A high level of CCoAOMT in ray parenchyma cells adjacent to living as well as dead vessels has been shown. These data therefore suggest that CCoAOMT is indeed involved in the synthesis of coniferyl alcohol units. In addition, the studies suggest that in living cells, the coniferyl alcohol is synthesised within the vessel and the adjacent living ray cells provide coniferyl alcohol to the adjacent vessel when it is dead. Monolignols are likely also provided by the adjacent xylem fiber cell; by immunolocalisation it was shown that CCoAOMT protein was present at the site of the cell that was adjacent to the vessel. The specific activity of the promoter in or adjacent the vessel can be used for several applications. It opens the possibility to reinforce vessel elements in transgenic plants that have altered fiber characteristics, allowing e.g. the construction of plants with fibers with a modified lignin content, without affecting the function of the vessels of the plant. Apart from lignin related products, the promoter system may be used to produce products, other than monolignols, that can be transported to the vessel. As a non limiting example, the promoter may be used to produce small antifungal peptides in the cells adjacent to the vessel. These peptides may be transported into the vessel and from there, to the whole plant, to confer resistance against fungal attack.

It has also been shown that both *Pop1* and *Pop2* promoters conferred expression in the periderm. The strong promoter activity in the periderm was associated with the formation of lignified cells in the phellem. Such observation suggests that CCoAOMT plays an important role in the generation of a protective dermal layer in plants. In addition, since the GUS activity was detected in the cell layer adjacent to the lignified layer, this supports the hypothesis that monolignols can be provided by adjacent cells.

In phloem, the promoter of *Pop2* confers expression within the phloem fiber cells, whereas the promoter of *Pop1* confers expression in the cells surrounding the phloem fibers. In xylem, however, both promoters are expressed very similarly. This suggests that *cis* elements necessary for xylem expression might be different from those needed for phloem expression. Hence, it might be possible to identify and

isolate elements that are needed specifically for expression associated with vessel elements. In the 4CL promoter, a negative *cis*-acting element which represses phloem expression has been identified. This element appears to be responsible for restricting vascular expression to the xylem (Hauffe et al., 1993).

Alignment of the two promoter sequences allowed us to identify putative elements involved in cell- and tissue-specific expression of these two genes. The AC-II element (or box L) present in the promoters of *Pop1* and *Pop2* had initially been identified by *in vivo* footprint experiments as light and elicitor-responsive promoter element in the parsley *PcPAL-1* and *Pc4CL* genes (Hauffe et al. 1993, Lois et al. 1989). They were later found to be present in various other genes of the phenylpropanoid pathway from other plants (Table 1). Da Costa e Silva (1993) isolated a DNA-binding factor BPF1 for the corresponding AC elements. They observed an induction of BPF1 synthesis in response to elicitor treatment in parsley cell suspension cultures. In spite of the convincing functional relevance, transient expression assays with promoter-reporter gene constructs in parsley were not sufficient to explain the responsiveness to light or elicitor, and a combinatorial interaction of these AC-elements even with exonic sequences has been suggested (Hauffe et al., 1993). The other homologous sequences, as identified in the consensus sequence (Fig. 13), or parts of it, optionally in combination with the AC-elements, may play a role in the determination of the tissue specificity of the promoter, especially in determining the expression in living vessel elements, in ray cells adjacent to vessels elements or in fibers adjacent to vessel elements.

Since the *CCoAOMT* promoters confer expression in cells adjacent to vessel elements but not next to fiber cells, it follows that specific signals must be transported from the vessel to the adjacent cells, even when the vessel element has undergone autolysis. In this respect, "cells, adjacent to vessel elements" are defined as those cells that can be reached by a signal molecule that is diffusing out of a vessel; cells that are not reached by the signal are considered as not adjacent. Said signal can also be a certain pressure originated from water that is transported through said vessel. Such a pressure can be captured by for instance a membrane protein which simply transports the signal concerned. For the expression of other lignification genes that are "constitutively" expressed in the living ray cells, such a signal would not a priori be necessary, because the expression of these genes could be part of the

default set of genes that are expressed in the ray cells. The presence of *cis* elements in the promoter of the *CCoAOMT* genes that interact with vessel derived factors can thus be anticipated. It is foreseen that these response elements are also located - at least partly - in the consensus sequence that is derived from the promoter comparison. These response elements, as part of the *CCoAOMT* promoter, or as separate elements in combination with another promoter sequence can be used to produce proteins in a reaction upon the signal coming from the vessel. As a non limiting example is mentioned that the signal is a toxic compound and the protein is a detoxifying enzyme, or the example where the signal is the result of a microbial contamination, and the protein an antimicrobial agent.

The close spatial association between promoter activity and lignified tissues strongly suggests the involvement of these two genes in the lignification process. Transgenic poplars in which the expression of *CCoAOMT* is reduced will unravel the precise role of *CCoAOMT* in lignification.

Table 1. Putative cis-acting elements and positions on various *PAL*, *4CL* and *CCoAOMT*

gene promoters

Promoter	Position	AC-element II	Reference
PvPAL2	-123	TCTCCACCAACCCC	Cramer et al. 1989
AtPAL1	-135	TCTCAACCAACTCC	Ohl et al. 1990
AtPAL2	-132	TCTCACCCACCCCT	Warner et al. 1995
PsPAL2	-204	TCTCAACCAACCAC	Yamada et al. 1994
PopPALg2b		TCTCAACCAACCCC	Osakabe et al. 1996
PopPALg1		TCTC ACCAACCAC	Osakabe et al. 1996
PcPAL1	-107	TCTC ACCTACCAA	Lois et al. 1989
PcPAL2	-107	TCTC ACCTACCAA	Logemann et al. 1995
PcPAL4	-268	TCTC ACCAACCCC	Logemann et al. 1995
Pc4Cl1	-128	TCTC ACCAACCCC	Logemann et al. 1995
Pc4Cl2	-128	TCTC ACCAACCCC	Logemann et al. 1995
St4Cl1	-17	TCTC ACCAACCAC	Joos et al. 1992
St4Cl2	-168	TCTC ACCAACCAC	Joos et al. 1992
PcCCoAOMT	-120	TCTC ACCAACCGC	Grimmig and Matern, 1997
Pop1(CCoAOMT)	-205 *	CCTC ACCAACCCC	this invention
Pop2(CCoAOMT)	-201 *	CCTC ACCAACCCC	this invention
	-498 *	CCTC ACCAACCCC	this invention

Consensus	TCTC	ACCAACCCC
	C	T AA

*At*, *Arabidopsis thaliana*; *Pc*, *Petroselinum crispum*; *St*, *Solanum tuberosum*; *Ps*, *Pisum sativum*; *Pv*, *Phaseolus vulgaris*; *Pop*, *Populus kitakamiensis* (Yuriko Osakabe), and *Populus trichocarpa* (this work). Position numbers counted from the transcriptional start except the ones marked with \* which were from the translation start codon.

Table 2. Promoter activity in transgenic poplar

	<u>CCoAOMT1-GUS</u>	<u>CCoAOMT2-GUS</u>
<u>Stem</u> :		
young internode	· xylem ray cells between vessels · parenchyma cells surrounding inner vessels	· xylem ray cells between vessels · parenchyma cells surrounding inner vessels
older internode	· xylem ray cells next to vessels * companion cells * parenchyma cells surrounding phloem fibres	· xylem ray cells next to vessels * phloem fibres
<u>Leaf and petiole</u> :	· xylem ray cells	· xylem ray cells * phloem fibres
<u>Root</u> :	· xylem ray cells next to vessels * phloem fibres	· xylem ray cells next to vessels * phloem fibres
<u>Periderm</u> :	· pellogen and phloclerm	· pellogen and phloclerm
<u>Bud</u>	· meristem	· meristem

Table 3. The deletions of CcoAOMT promoters activity in stems of transgenic poplar

	number of analysed primary transformants	primary tissue	secondary tissue	response to wounding
pBINPOP1 (- 1993)	8(15)	primary xylem, base of hair	vessel, ray cells next to vessel, periderm	yes
pBIN1DA5 (-456)	5(6)	pith ray cell surrounding inner of xylem primary xylem, base of hair	companion cell vessel, ray cells next to vessel, periderm	yes
pBIN1DB3 (-199)	13(13)	pith ray cell surrounding inner of xylem undetected	ray initial cells in cambial zone undetected	no
pBIN1DC2 (-184)	10(10)	undetected	phloem fibres, cortex and periderm	no
pBIN1DD1 (-114)	8(8)	undetected	undetected	no
pBINPOP2 (- 1362)	8(15)	primary xylem, base of hair	vessel, ray cells next to vessel, periderm	yes
pBIN2DA4 (-497)	4(7)	pith ray cell surrounding inner of xylem primary xylem, base of hair	phloem fibres vessel, ray cells next to vessel, periderm	yes
pBIN2DB3 (-195)	12(12)	pith ray cell surrounding inner of xylem undetected	phloem fibres undetected	no
pBIN2DD1 (-110)	10(10)	undetected	undetected	no

## References

- Albrecht, K.A., Wedin, W.F. and Buxton, D.R. (1987). Cell-wall composition and digestibility of alfalfa stems and leaves. *Crop Sci.* 27, 735-741.
- Atanassova, R., Favet, N., Martz, F., Chabbert, B., Tollier, M.-T., Monties, B., Fritig, B. and Legrand, M. (1995) Altered lignin composition in transgenic tobacco expressing O-methyltransferase sequences in sense and antisense orientation. *Plant J.* 8, 465-477.
- Bevan, M., Shufflebottom, D., Edwards, K., Jefferson, R., and Schuch, W. (1989). Tissue- and cell-specific activity of a phenylalanin ammonia-lyase promoter in transgenic plants. *EMBO J.* 8, 1899-1906.
- Bevan, M. W. (1984). Binary Agrobacterium vector for plant transformation. *Nucl. Acids Res.* 12, 8711-8721.
- Boudet, A.M., Lapierre, C. and Grima-Pettenati, J. (1995) Biochemistry and molecular biology of lignification. *New Phytol.* 129, 203-236.
- Bugos, R.C., Chiang, V.L.C., and Campbell, W.H. (1991). cDNA cloning, sequence analysis and seasonal expression of lignin-bispecific caffeic acid/5-hydroxyferulic acid O-methyltransferase of aspen. *Plant Mol. Bio.* 17, 1203-1215.
- Buxton, D.R. and Russell, J.R. (1988) Lignin constituents and cell-wall digestibility of grass and legume stems. *Crop Sci.* 28, 553-558.
- Casler, M.D. (1987). In vitro digestibility of dry matter and cell-wall constituents of smooth brome grass forage. *Crop Sci.* 27, 87-99.
- Chapple, C.C.S., Vogt, T., Ellis, B.E. and Somerville, C.R. (1992). An Arabidopsis mutant defective in the general phenylpropanoid pathway. *Plant cell* 4, 1413-1424.
- Collazo, P., Montoliu, L., Puigdomènech, P. and Rigau, J. (1992). Structure and expression of the lignin O-methyltransferase gene from *Zea mays* L. *Plant Mol. Biol.* 29, 857-867.

Cramer, C.L., Edwards, K., Dron, M., Liang, X., Dildine, S., Bolwell, G.P., Dixon, R.A., Lamb, C.J. and Schuch, W. (1989) Phenylalanine ammonia-lyase gene organisation and structure. *Plant Mol. Biol.* 12, 367-383.

Da Costa e Silva, O., Klein, L., Schmelzer, E., Trezzini, G.F. and Hahlbroeck, K. (1993) BPF-1, a pathogen-induced DNA-binding protein involved in the plant defense response. *Plant J.* 4, 125-135.

Davin, L.B., and Lewis, N.G. (1992). Phenylpropanoid metabolism: biosynthesis of monolignols, lignans and neolignans, lignins and suberins. In *Phenolic Metabolism in Plants*, H.A. Stafford and R.K. Ibrahim, eds (New York: Plenum Press), pp. 325-375.

Dixon R.A., Maxwell, C.A., Ni, W., Oommen, A. and Paiva, N.L. (1994) Genetic manipulation of lignin and phenylpropanoid compounds involved in interaction with micro-organisms. *Rec. Adv. Phytochem.* 28, 153-178.

Feuillet, C., Lauvergeat, V., Deswarte, C., Pilate, G., Boudet, A., and Grima-Pettenati, J. (1995). Tissue- and cell-specific expression of a cinnamyl alcohol dehydrogenase promoter in transgenic poplar plants. *Plant Mol. Biol.* 27, 651-667.

Gowri, G., Bugos, R.C., Campbell, W.H., Maxwell, C.A., and Dixon, R.A. (1991). Stress responses in alfalfa (*Medicago sativa* L.). X. Molecular cloning and expression of S-adenosyl-L-methionine: caffeic acid 3-O-methyltransferase, a key enzyme of lignin biosynthesis. *Plant Physiol.* 97, 7-14.

Grimmig, B. and Matern, U. (1997). Structure of the parsley caffeoyl-CoA O-methyltransferase gene, harbouring a novel elicitor responsive *cis*-acting element. *Plant Mol. Biol.* 33, 323-341.

Grisebach, H. (1981) Lignins. *The Biochemistry of Plants*. Vol. 7. E.E. Conn, ed (New York: Academic Press), pp. 457-478.

Haufler, K.D., Lee, S.P., Subramaniam, R., and Douglas, C.J. (1993). Combinatorial interactions between positive and negative *cis*-acting elements control spatial patterns of *4Cl-1* expression in transgenic tobacco. *Plant J.* 4, 235-253.



Hauffe, K.D., Paszkowski, U., Schulze-Lefeert, P., Hahlbrock, K., Dangl, J.L., and Douglas, C.J. (1991). A parsley 4Cl-1 promoter fragment specifies complex expression patterns in transgenic tobacco. *Plant cell* 3, 435-443.

Hawkins, S., Samaj, J., Lamergeat, V.A., and Grima-Pettenati, J. (1997) Cinnamyl Alcohol Dehydrogenase: Identification of New Site of Promoter Activity in Transgenic Poplar. *Plant Physiol.* 113: 321-325.

Jefferson R.A., Kavanagh T.A., and Benan M.W. (1987). GUS fusions:  $\beta$ -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* 6, 3901-3907.

Joos, H.J. and Hahlbrock, K. (1992) Phenylalanine ammonia-lyase in potato (*Solanum tuberosum* L.). Genomic complexity, structural comparison of two selected genes and modes of expression. *Eur. J. Biochem.* 204, 621-629.

Jung, H.G. and Vogel, K.P. (1986). Influence of lignin on digestibility of forage cell wall material. *J. Anim. Sci.* 62, 1703-1712.

Kneusel R.E., Matern U. and Nicolay K. (1989). Formation of trans-caffeoyl-CoA from trans-4-coumaroyl-CoA by  $Zn^{++}$  dependant enzymes in cultured plant cells and its activation by an elicitor induced pH shift. *Archives of Biochemistry and Biophysics.* 269:455-462.

Kühnl T., Koch U., Haller W., Wellmann E. (1989) Elicitor induced S-adenosyl-L-methionine : caffeoyl-CoA 3-O-methyltransferase from carrot cell suspension cultures. *Plant science* 60, 21-25.

Leplé, J.C., Brasileiro, A.C.M., Michel, M.F., Delmotte, F., and Jouanin, L. (1992). Transgenic poplars: expression of chimeric genes using four different constructs. *Plant Cell Rep.* 11, 137-141.

Lewis, N.G., and Yamamoto, E. (1990) Lignin: Occurrence, biogenesis and biodegradation. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 41,455-496.

Logemann, E., Parniske, M. And Hahlbrock, K. (1995) Modes of expression and common structural features of the complete phenylalanine ammonia-lyase gene family in parsley.

*Proc. Natl. Acad. Sci. USA* 92, 5905-5909.

Lois, R., Dietrich, A., Hahlbrock, K. and Schulz, W. (1989) A phenylalanine ammonia-lyase gene from parsley: structure, regulation and identification of elicitor and light responsive cis-acting elements, *EMBO J*, 8, 1641-1648

Matern U, and Grimmig B (1994) Natural phenols as stress metabolites. In: Geibel M, Treutter D, Feucht W (eds) *Acta Horticulturae: International Symposium on Natural Phenols in Plant resistance*. pp. 448-462. Sellier Druck, Freising, Germany.

Neish A.C. (1968) Monomeric intermediates in the biosynthesis of lignin. In : Constitution and biosynthesis of lignin. K. freudenberg and A.C. Neish, eds (new York: Springer-Verlag):3-43.

Nicholson, R.L. and Hammerschmidt, R. (1992) Phenolic compounds and their role in disease resistance. *Annu Rev. Phytopath* 30: 369-389 .

Ohl, S., Hedrick, S.A., Chory, J. And Lamb, C.J. (1990) Functional properties of a phenylalanine ammonia-lyase promoter in *Arabidopsis*. *Plant Cell* 2, 837-848.

Osakabe-Y, et al. (1996). Immunocytochemical localization of phenylalanine ammonia-lyase in tissues of *Populus kitakamiensis*. *Planta* 200(1): 13-19.

Peleman, J., Boerjan, W., Engler, G., Seurinck, J., Botterman, J., Alliotte, T., Van Montagu, M., and Inze, D. (1989) Strong cellular preference in the expression of a housekeeping gene of *Arabidopsis thaliana* encoding *S-adenosylmethionine synthetase*. *Plant cell* 1, 81-93.

Saka, S., and Goring, D.A.I. (1985). Localization of lignins in wood cell walls. In *Biosynthesis and Biodegradation of Wood Components*, T. Higuchi (ed.). Orlando, Academic Press, pp. 51-62.

Shure, M., Wessler, S. And Federoff, N. (1983) Molecular identification and isolation of the Waxy locus in maize. *Cell* 35, 225-233.

Smith C.G., Rodgers M.W., Zimmerlin A., Ferdinando D., Bolwell G.P. (1994). Tissue and subcellular immunolocalisation of enzymes of lignin synthesis in differentiating and wounded

hypocotyl tissue of French bean (*Phaseolus vulgaris* L.). *Planta* 192: 155-164.

Speer, E.O. (1987). A method of retaining phloroglucinol proof of lignin. *Stain Technol.* 62, 279-280.

Tien, M. (1987) Properties of ligninases from *Phanerochaete chrysosporium* and their possible applications. *CRC Crit. Rev. Microbiol.* 15, 141-168.

Yamada, T., Sriprasertsak, P., Kato, H., Hashimoto, T., Shimizu, H. And Shiraishi, T. (1994) Functional analysis of the promoters of phenylalanine ammonia-lyase genes in pea. *Plant Cell Physiol.* 35, 917-926.

Ye, Z.-H. and Varner, J.E. (1995) Differential expression of two O-methyltransferases in lignin biosynthesis in *Zinnia elegans*. *Plant Physiol.* 108, 459-467.

Ye, Z.-H. Kneusel, R.E., Matern, U. and Varner, J.E. (1994) An alternative methylation pathway in lignin biosynthesis in *Zinnia*. *Plant Cell* 6, 1427-1439.

Van Doorselaere, J., Baucher, M., Chognot, E., Chabbert, B., Tollier, M.-T., Petit-Conil, M., Inze, D., Boerjan, W. and Jouanin, L. (1995) A novel lignin in poplar trees with a reduced caffeic acid/ 5-hydroxyferulic acid O-methyltransferase activity. *Plant J.* 8, 855-864.

Zahm et al.(1984) Site specific mutagenesis of the Ti plasmid by transformation of *Agrobacterium tumefaciens* with mutagenized T-DNA fragments cloned in *E. Coli* plasmids. *Mol. Gen. Genet.* 194, 188-194.

Claims

1. An isolated DNA sequence according to figure 1 B (SEQ.ID.NO.1) representing a *CCoAOMT* promoter or a functional part thereof having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel.
2. An isolated DNA sequence according to figure 1 C (SEQ.ID.NO.2) representing a *CCoAOMT* promoter or a functional part thereof having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel.
3. An isolated DNA sequence according to claim 1 or 2 representing a *CCoAOMT* promoter or a functional part thereof having said biological activity wherein the cells adjacent to the vessel are xylem ray cells.
4. An isolated DNA sequence according to claim 1, 2 or 3 representing a *CCoAOMT* promoter or a functional part thereof having said biological activity wherein the vessel is a differentiating vessel.
5. A promoter sequence conferring expression in living vessel elements and/or in ray cells, adjacent to living or autolysed vessel elements and/or in fibers adjacent to said living or autolysed vessel elements, with the proviso that said expression does not occur constitutively in xylem fiber cells or ray cells that are not adjacent to said vessel elements.
6. An isolated DNA sequence according to claim 1 wherein the functional part represents the region from -114 to - 456, preferably from -199 to - 456 according to the numbering indicated in figure 1B.
7. An isolated DNA sequence according to claim 2 wherein the functional part represents the region from - 110 to - 497, preferably from - 195 to - 497 according to the numbering indicated in figure 1C.
8. A recombinant DNA comprising any of the isolated DNA sequences according to claim 1 and/or 2.
9. A plant or tree cell comprising a recombinant DNA according to claim 8 integrated

in its genome.

10. A transgenic plant or tree comprising the cell according to claim 9 and its progeny thereof.




Figure 1




Figure 1  
Continued






Figure 1. The lignin biosynthesis pathway.

PAL, phenylalanine ammonia-lyase; C4H, cinnamate-4-hydroxylase; C3H, coumarate-3-hydroxylase; COMT, bispecific caffeic acid/5-OH-ferulic acid-*O*-methyltransferase; F5H, ferulate-5-hydroxylase; CCoA3H, coumaroyl-CoA-3-hydroxylase; CCoAOMT, caffeoyl-CoA-*O*-methyltransferase; 4CL, 4-OH-cinnamate-CoA ligase; CCR, cinnamoyl-CoA-reductase; CAD, cinnamyl alcohol dehydrogenase.




Figure 1-A. Structure of CCoAOMT genes.  
A: CCoAOMT gene1, *pop1*. B: CCoAOMT gene2, *pop2*.  
C: cDNA.

BNSDOCID: &lt;WO\_9909188A2\_1\_&gt;

[illegible]

BNSDOCID: <WO\_\_\_9909188A2\_I\_>

1130 1150 1170 1190 1210 1230  
ACAGGGACTTTGTTTGGAGCTCAACAAGGCACCTTCTGCTGACCCGAGGATTGAAATTTGGATGCTTCTGTTGGTGATGGCATTCACTCTCTCCGTCGGATCCAATgaggagacctgcc  
1250 1270 1290 1310 1330 1350  
agtattgttatctgagtggaccattgaaatgggtcactttacaagaacaaggagatgcaaatagttgtttttaccacactttgattttcaatgggttataatttgttacttgaacagaatg  
1370 1390 1410 1430 1450 1470  
gtgtatgattgagaaattcctctcttaaaattctgtgaagtggattttttatgcacttaataatattgttcgggtggctaaacacttggtagttgtatgcattgctaagatggagatttc  
↓ Poly A cleavage site  
1490 1510 1530 1550 1570 1590  
tcattctatcccaaggccatcatagtttaaccagtttacaactaaattctcgagaaaagggtttgttcccaattaaagtgttcttagacattatgaatgattgtrataaaaatggttccaaaaacttct  
1610 1630 1650 1670 1690 1710  
aatccgttggacttcttttttggtaaaaatgttttttatgttttccaaagtgtttttgttagacggtgagaaaaacaagaagcgtgtacgagtgtacctactagtgtgctaactagtcacttttag  
1730 1750 1770 1790  
agtttagagacagacattgaggttggcggaggaataatccgaaattatcttgaacaagtcataacttttcacacgaat

Figure 1-C. DNA sequence of *PtCCoAOMT2* (AJ223620), from *P. trichocarpa* cv trichobol. The coding regions are indicated as capital letters. The TATA-box and AC-II elements are underlined. The primers used to generate the deletions of the promoter are indicated in *italics*.

-1360	-1340	-1320	-1300	-1280	-1260
tcagagaaacacggttccacacggttcccaaacatgatttaatatatcacggttatattttgcatttcaatagggttttgaaaaaagaatgaattttattttattttgttttaaa					
-1240	-1220	-1200	-1180	-1160	-1140
ttatttttttagtatttttagatcgttttaacatgccaatgtccaaaaaatttaaaaaaatattatttcaattttatttggtaagaaaaaaacctcaaaaaacaattattaccaaaaacacac					
-1120	-1100	-1080	-1060	-1040	-1020
aaactggccttgaagtaacctatagcataatgcactaaccaattattttaattttccatccctgcctcatggagaagattccatggttgaagacigtatgataaggaaaaaatgtcatgaac					
-1000	-980	-960	-940	-920	-900
tcarggtataagtaatttccratccataaacagcaagcttgatgttaggttgatgtttcttcttcatggaaatgttttgccatgccacacgaacaggcaagaaaaaacagg					
-880	-860	-840	-820	-800	-780
acaaattataggaaatgtctcaatgcatgtatattaaadatatatttttgaatttgaaaaattatttttaatatattataaataaaaaatacttaaaaaaacatcttttgcactttaa					
-760	-740	-720	-700	-680	-660
aaaaacatttaattaatctgcctagcttttactagaaaaacacacactaaactgggcgtatgaaaactccagccatttttatatatattgtccctggtattatcatagacgtaaaaacgaaat					
-640	-620	-600	-580	-560	-540
tggatttttttaattttgttggagaaaaaaaagaaaaataataattgtccagcagtaagaaggagagattctttaaaaaggagtcacattgtcaatgcggttgctacagagcccaactcc					
-520	-500	-480	-460	-440	-420
cgtagagtcgaattcttggaggacacctccacacaccccttaccacactttctattagcagcacatgtagccatcccccaacaacaagtggtgagccccaccacaatttttctactctctacgat					
-400	-380	-360	-340	-320	-300
ttaaatcaatttacacgtggcatataaatgtcagacctttttatttcaagaaaccaaactaacaccgtagaacttaatttttttcgcaaatcttaaaaataagggtacatgaattaaatgat					
-280	-260	-240	-220	-200	-180
agaaattgaattagtgtccgaaacctcaaaatgaccactggacaaacacgataagtgggtcccaaaaatacccacggtgtcttaagaaactcaccccaacccacccggttgggaagcgggtcc					
AC-II element					

10/36

-160  
 aaccacccactactccgggtcaaaaccgggactctcatctcccaataaataaccacccgcccctggccatttccaatcagggtcagacatccttaaccatcg:cgcgccccagaaaaaccttccaac  
 -140  
 TATA  
 -120  
 -100  
 -80  
 -60  
 Transcription start site  
 -40  
 -20  
 0  
 20  
 40  
 60  
 gccaggaaagagagatagttttgttataagatafacaataaataATGGCCGCCAACGGAGAGAGAAAGCAGACTCAGCCGGGAGGCATCAAGAAAGTTGGCCACAAAGAGCCTTTTGTCAAAG  
 Translation start site  
 80  
 100  
 120  
 140  
 160  
 180  
 TGATGCTCTTTACCAAGtaatttaaccggagaaacccctgatcttgggtgcgagttttgttttttttttcccttggtttgttttttttaacattttttgtatatatttggattgggttttttag  
 200  
 220  
 240  
 260  
 280  
 300  
 TATATTCTTGAGACCAAGTGTATACCCAGAGAGCCTGGAATGCATGCAAGAGGCTTAGAGAGTTGACTGCCAAGCATCCTTGGtatgtttgttgaaatcctccatgcatttttaaatacatca  
 320  
 340  
 360  
 380  
 400  
 420  
 acatgagagatttttttccataaaaaaaaggggttttttttttctatatattgatgttttatggcgtggttttaataaatctgatgtgaatgatgggaaacagGAACATCATGACCA  
 440  
 460  
 480  
 500  
 520  
 540  
 CATCTGCTGATGAGGSCAATCTTGAAACATGCTTTTAAAGCTTATCAATGGCAAGAACACCATGAGAGATTGGTGTGTTTCACTGGCTATCTCTCTTGGCCACTGCTCTTGGTATATCCCTG  
 560  
 580  
 600  
 620  
 640  
 660  
 AGGATGGAAAGtaaaactaaagacctgagattttctctgtctcaaaacagtcuaaagaaatttaggtgacgtaagaagaaataattgggaatctcttgaaattgcagATCTTGGCTATGGACH  
 680  
 700  
 720  
 740  
 760  
 780  
 TCACACAGAAAACCTATGAACCTGGCTCTCCCGGTGATTCAGAAAGCTGCTCTGGACACACAGATTGAGTTCAAGGAAGGCCCTGCTTGGCACTTCTGGATCAAAATGATTGAAGATgtaa  
 800  
 820  
 840  
 860  
 880  
 900  
 gaatatattctgtctttgacuaaaaaaaacattttttgtttgtctgtatgagaaagagatgatattcatttaagaaaaattgatgtgctattacttagcttactattctgtgtgtggcg  
 920  
 940  
 960  
 980  
 1000  
 1020  
 aactacagggaaagTACCATTGACTTATGACTTTCATCTTTTGGATGCTGTACAGAGGCAATATATTAACCTACACAGAGGTTGATTGAGCTTGTCAAGTGGGAGGGTGTGATTGGGT  
 1040  
 1060  
 1080  
 1100  
 1120  
 1140  
 ATGACACACCTGTGGGATGGATCTGTGGTGGCACCGGATGGCCCAATGAGGAATATGTGAGGTACTATAGGGACTTTGTTCTGGAGCTTCATAGGACACTTGCACCTGACCCCA  
 1160  
 1180  
 1200  
 1220  
 1240  
 1260  
 GGATTGAATCTGCATGCTTCCTGTGTGGTGATGATCACTCTCTGGCGGTGCGATCAAGTgaggggctgcattccctgcgatattatatatatatacaaatgttgaccattgatttggctc

SUBSTITUTE SHEET (RULE 26)

1280 1300 1320 1340 1360 1380  
acttgcacaaacagtgatggtgaataaagagattttaccgaccttgcctatatatttgacctgaagaatggcgaatggccgagaaactccattttgaatttctgtctaaatggatt  
Poly A signal  
1400 1420 1440 1460 1480 1500  
ttttacgcacttaatccatattgtttctttaaccaagtaacttatcacgcgggtgctgagaatgttcatggtttaaggagtttgccttcaacattcaaaaattaaagaagcacgcgggtcccttgct  
↓ Poly A cleavage site  
1520 1540 1560 1580 1600 1620  
cccttacgaagaccactagatgcggttcccatcagtttaatgaccttaaggacgagtttatctcttctgctgactttctgtacatgcagccgcactctcctcglaataaataaact  
1640 1660 1680 1700  
gctacccgcaatgttaaaatgattgtgtatcgagaaagggttttttcuagcttgccttgaactagcgtggaaaa



1151  
 prom1-2.msf{pccoamt2} TT TTTATATA TTTGTCCTGT GATTATCAT. ....  
 prom1-2.msf{pccoamt1} TATATGCATA CCAAGTATAC CCTTTACATC TGCCCTAAAT AGACGGTAA  
 Consensus T -T-T- ATA ----- T--- TT-CAT-----  
 -AGAC- GTAA  
 Upper primer

1201  
 prom1-2.msf{pccoamt2} AACGAAATTG GATTTTT ..... TT ATTTGTTGG AGAAAAAAA  
 prom1-2.msf{pccoamt1} AACGGATTG GATTTTT GCC GACAAAGGCT AGTTTG.TGG AGAAAAAACA  
 Consensus AACG-A- TTG GATTTTT----- -T A- TTG- TGG AGAAAAAA -A

1251  
 prom1-2.msf{pccoamt2} AAGAAAAATAA ATATTGTCAG CAGTAAGAC. ....  
 prom1-2.msf{pccoamt1} ACGAAGATAA ATTTTAGAT GACAAAGTCA ACAATAGTTC  
 Consensus A -GAA -ATAA AT -TT--- A- -A- -AAG-C----- -GAGAGATTCT

1301  
 prom1-2.msf{pccoamt2} TAAAGGAGT CATCCATTGTC CAATGCGGTG GCTACGAGCC ACCAACTCC  
 prom1-2.msf{pccoamt1} TTAAAGGACT CATCCGTTG. ACGGAGGTG GCCATATGCT ACCAACTCT  
 Consensus T- AAAGGA- T CATCC- TTG- - A- -G- GGTG GC- A--- GC- ACCAACTC--

1351  
 prom1-2.msf{pccoamt2} GTGGAGTCAA ATTCTTGAGG ACACCTCACC AACCCCTTAC CCACCTTCTA  
 prom1-2.msf{pccoamt1} G .GACGTGGA GTCCCTTTGG TAATTTCACC TATCCCTCAC CCAATTCTA  
 Consensus G- G-- GT- A - T- C- T-- GG- -A-- TCACC -A-CCCT-AC CCA- TTTCTA

1401  
 prom1-2.msf{pccoamt2} TTAGC... A GCACATGTA ..... GCCATCCCCA ACAACAAAGT  
 prom1-2.msf{pccoamt1} TTAGCAGTTA GCACATGTA TTTATGATTG GTGAGCCAG CACAAATCTT  
 Consensus TTAGC--- -A GCACATGTA----- G-- A- CCC----- A-A--- T

Fig 1-d

1451  
prom1-2.msf{pccoaomt2} GG TGAGCCCA CCACA.....  
prom1-2.msf{pccoaomt1} TT CCAGTTAA ACACATATAT TAATTATGA TTAATTATTT AATTCTCTCC  
Consensus - ---AG- --A- CACA-----  
AATT -T-CTCTC-

1501  
prom1-2.msf{pccoaomt2} ACGATT. TA AATCAATTAC ACGTGGCATA AATGTCGAG CCTTTTATTT  
prom1-2.msf{pccoaomt1} ACTCTTAACA AATTAAATCAT ACATGGCATA ACAT TTIAG CTTTIGATCT  
Consensus AC --TT--- A AAT-AAT-A- AC-TGGCATA A- AT- T- -AGC-TTT-AT-T

1551  
prom1-2.msf{pccoaomt2} CAAGAAACCA AACCTAACAC CGTGAACTTA ATTCT. TTC GCAAATATCT  
prom1-2.msf{pccoaomt1} CGAGAAATCTC TACCTA.. AC CATTGACTTC TTTACTGTTT AGGAATCTTA  
Consensus C -AGAA-C-- -ACCTA- -ACC-T-- ACTT- -TT CT- TTC --- AAT -T--

1601  
prom1-2.msf{pccoaomt2} AGAAATTGAA TTAGTGTG.. CGAAACCTAA AATGAC.....  
prom1-2.msf{pccoaomt1} GAACCGATAT TTGGTGTGTCAT AGATCCCAAA AATGACGCCA GCGATGCCCTA  
Consensus -A---- -A- TT- GTGTC-- -GA --CC -AA AATGAC-----

1701  
prom1-2.msf{pccoaomt2} .....  
prom1-2.msf{pccoaomt1} AGGGAAGGAG TACCACTAGC CCACAGCAG ATACG ATCAC CAACAAGGTG  
Consensus -----CAC-- -AC -A-CAC C- A-AA -GTG

1751  
prom1-2.msf{pccoaomt2} GGTCCCA.....  
prom1-2.msf{pccoaomt1} GGTCCCAAT TTGGTGGGCC AAAAACCAC ATTATCCTTC GTCCTAACTA  
Consensus GGTCCCA-----  
AAA-ACCCAC- T- TCCT-- G-----





Figure 1-D. Comparison of DNA sequences of the two promoters of CCoAOMT genes from poplar. The upper line, pccoamt2, is promoter of pop2. The lower line, pccoamt1, is promoter of pop1. The part of coding sequence is in bold type. Conserved promoter regions are indicated by open boxes. The AC-element II are marked underline. The sequence represented by horizontal arrows are chosen as primers for amplifying promoters from different species in poplar. The vertical arrow is indicated translation start site.

<End of Fig. 1-D>



15/36

**Fig. 2. Structure of expression vectors**

PTB1: chimaeric CCoAOMT class I promoter-gus cassette.

PTB2: chimaeric CCoAOMT class II promoter-gus cassette.




Fig. 3.

Histochemical localization of GUS activity in transgenic plant

The cross section of young internode were double-stained. Lignified cells and tissues are revealed by red color after phloroglucinol-HCl staining. GUS activity is only detected in the xylem ray cells located between the lignified vessels.

*Amorphous dia-*

Fig • 4. Histochemical localization of GUS activity in transgenic poplar. The transverse section through young stem shown GUS activity localized in vessels and in the ray cells surrounding and between vessels.





Fig • 5. Histochemical localization of GUS activity in transgenic poplar stems. The transverse sections shown GUS activity was localized in vessels and in the ray cells adjacent to vessels in xylem tissue of old stems: A, in pBINPOP1 transform; B, in pBINPOP2 transform.

A.



B.





Fig • 6. Dark-field image of transverse sections through stem in transgenic poplar.  
The red-purple represents GUS activity.










Fig. 7. Histochemical localization of GUS activity in transgenic poplar stems. A, the section was stained with phloroglucinol and GUS, lignified walls stain red. B, GUS activity localization with CCoAOMT2-GUS construct. C, GUS activity localization with CCoAOMT1-GUS construct.

Fig. 8. Enlargement of transverse sections of stem shown GUS activity: A, in the companion cells; B, in the phloem fibres; and C, in periderm.


A.




B.



C.



22/36



**Fig. 9:**  
**Histochemical localization of GUS activity in transgenic poplar petioles.**  
**A:** The section was stained with phloroglucinol; lignified walls stain red.  
**B:** GUS activity localization with *CCoAOMT2*-GUS construct.  
**C:** GUS activity localization with *CCoAOMT1*-GUS construct.




Fig. 10


**Histochemical localization of GUS activity in transgenic poplar leaves.**

**A: The section was stained with phloroglucinol; lignified walls stain red.**

**B: GUS activity localization with CCoAOMT2-GUS construct.**

**C: GUS activity localization with CCoAOMT1-GUS construct.**

Fig. 11. GUS staining in transgenic poplar.  
A and B, in root; C, in meristem of shoot bud; D, in leaf.






Fig. 12. Immunolocalization of CCoAOMT in the stems of *P. trimula* x *P. alba*. CCoAOMT was localized intensively in the xylem ray cells adjacent to vessels and in the phloem fibres (C). A, control reaction.

BNSDOCID: &lt;WO\_\_9909188A2\_I\_&gt;

pgdel: *P. deltooides* S 9-2 B IBW  
pgalb: *P. alba*  
pgcil: *P. ciliata* 72085 I  
pgnig: *P. nigra italica* aral  
pgmig: *P. nigra* Ghoy B IBW  
pbdel: *P. deltooides* S 336-4 B IBW  
pgeup: *P. euphratica*;  
pglau: *P. laurifolia* 69/65  
pgen: *P. trichocarpa* vc trichobel (this invention)  
number 1 and 2: refer to PtCCoAOMT1 and PtCCoAOMT2, respectively  
a, b and c: different alleles.  
1D1: **pBIN1DD1**; 1D2: **pBIN1DC2**; 1D3: **pBIN1DB3**; 1D5: **pBIN1DA5**; 2D1: **pBIN2DD1**; 2D3: **pBIN2DB3**; 2D4: **pBIN2DA4**

BNSDOCID: &lt;WO 9909188A2\_I\_&gt;



BNSDOCID: <WO\_\_9909188A2\_1\_>

p1eup_49.msf(pgeuplcrev)	cgaagaacct.	.....ct	Accta..acc	at.....aa	acct..attta	ttgttctaga	atcttagaaa	aaaggaggac	..aaaaaaa
p1eup_49.msf(pgeuplarev)	cgaagaacct.	.....ct	Accta..acc	at.....aa	acct..attta	ttgttctaga	atcttagaaa	aaaggaggac	..aaaaaaa
p1eup_49.msf(pgeuplbreve)	cgaagaacct.	.....CT	accta..acc	at.....aa	acct..attta	ttgttctaga	atcttagaaa	AAAggaggAC	..aaaaaaa
p1eup_49.msf(pgalbic)	cgaagaacct.	.....CT	accta..acc	atgaacaata	acct..aatta	ttgttcaaga	atcttagaaa	aaaggaggac	..aaaaaaa
p1eup_49.msf(pgalbia)	cgaagaAtct.	.....CT	accta..acc	atgaacaata	acct..aatta	ttgttcaaga	atcttagaaa	aaaggaggac	..aaaaaaa
p1eup_49.msf(pgnigia)	cgaataacct.	.....CT	acccta..acc	atg.....a	tctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pbmabib)	cgaataacct.	.....CT	acccta..acc	atg.....a	tctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pbnigib)	cgaataacct.	.....CT	acccta..acc	atg.....a	tctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pbmigib)	cgaataacct.	.....CT	acccta..acc	atg.....a	tctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pgcilic)	cgaagaacct.	.....CT	accta..acc	atg.....a	gctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pgcilb)	cgaagaacct.	.....CT	accta..acc	atg.....a	gctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pglaua)	cgaagaacct.	.....CT	accta..acc	atg.....a	gctt..attta	ttgttcaaga	atcttagaaa	aaaggaggaggg	..ggaataaaa..
p1eup_49.msf(pbmabib)	cgaagaatctt	agaaaaaagg	aggacaaatt	accta..acc	atg.....a	acct..attta	ctgttcaaga	atcttagaaa	aaaggaggaggg
p1eup_49.msf(pbdellb)	cgaagaatctt	agaaaaaagg	aggacaaatt	accta..acc	atg.....a	acct..attta	ctgttcaaga	atcttagaaa	aaaggaggaggg
p1eup_49.msf(pbdellb)	cgaagaatctt	agaaaaaagg	aggacaaatt	accta..acc	atg.....a	acct..attta	ctgttcaaga	atcttagaaa	aaaggaggaggg
p1eup_49.msf(pccoaoom1)	cgaagaatct.	.....CT	accta..acc	att.....g	acct..cttta	ctgttcaaga	atcttagaaa	aaaggaggaggg	..aaaaaaa..a
p1eup_49.msf(pgeup2b)	taatatctca	a.....	acctaacacc	gtg.....a	acttaatttc	tttcgcaaaa	acct.....aaa	ataagggggac	ggaaatttaa
p1eup_49.msf(pgeup2arev)	taatatctca	a.....	acctaacacc	gTT.....A	acttaatttc	ctttcaaaa	atct.....aaA	ataagggggac	ggaaatttaa
p1eup_49.msf(pgdell2b)	cagaagaacca	g.....	acctaacacc	gtg.....a	acttaatttc	tttcgcaaat	atct.....aaa	ataagggggac	atgaatt....
p1eup_49.msf(pgdell2a)	cagaagaacca	g.....	acctaacacc	gtg.....a	acttaatttc	tttcgcaaat	atct.....aaa	ataagggggac	atgaatt....
p1eup_49.msf(pccoaoom2)	cagaagaacca	a.....	acctaacacc	gtg.....a	acttaatttc	tttcgcaaat	atct.....aaa	ataagggttac	atgaatt....
Consensus	--A-A--C--	-----	ACCTA--ACC	AT-----	-C-----TT-	-T-----	A-CT-----	AA	A-A--G-----AA-----

**SUBSTITUTE SHEET (RULE 26)**

pileup_49.msf(pbmmax1a)	acccctccc	ggttgtagc	cgggtcagc	tcccgcctac	tccaattcaa	acggggctct	catttccaat	aaataccacc	cgcctttac	cattttcgat
pileup_49.msf(pbdell1b)	acccctccc	ggttgtagc	cgggtcagc	tcccgcctac	tccaattcaa	acggggctct	catttccaat	aaataccacc	cgcctttac	cattttcgat
pileup_49.msf(pbdell1a)	acccctccc	ggttgtagc	cgggtcagc	tcccgcctac	tccaattcaa	acggggctct	catttccaat	aaataccacc	cgcctttac	cattttcgat
pileup_49.msf(pccoaom1)	acccctccc	ggttgtagc	cgggtcagc	tcccgcctac	tccaattcaa	acggggctct	catttccaat	aaataccacc	cgcctttac	cattttcgat
pileup_49.msf(pgeup2b)	accccaacc	ggttgtagc	cgggtcagc	accccaacc	tccggtcaca	acgggaactct	cgttccaat	aaataccacc	tgccctt.gc	cattttcaat
pileup_49.msf(pgeup2arev)	accccaacc	ggttgtagc	cgggtcagc	accccaacc	tccggtcaca	acgggaactct	cgttccaat	aaataccacc	tgccctt.gc	cattttcaat
pileup_49.msf(pgdell2b)	accccaacc	ggttgtagc	cgggtcagc	accccaacc	tccggtcaca	acgggaactct	cgttccaat	aaataccacc	tgccctt.gc	cattttcaat
pileup_49.msf(pgdell2a)	accccaacc	ggttgtagc	cgggtcagc	accccaacc	tccggtcaca	acgggaactct	cgttccaat	aaataccacc	tgccctt.gc	cattttcaat
pileup_49.msf(pccoaom2)	ACCC--CCC	GTTGG-AGC	CGGTCCAGC	TCCCG-TAC	TCCAATCAA	ACCGGGCTCT	CATTTCAAI	AAATA-CACC	-GCCCTT--C	CA-TTT--A-
Consensus		T	A	A	A	G	G	G	T	A
		↑ D2 (oligo 19)								
pileup_49.msf(pgeup1crev)	caggt.....	.....	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgeup1arev)	caggt.....	.....	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgeup1brev)	caggt.....	.....	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgalbic)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgalb1a)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgnig1a)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pbmmax1b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgnig1b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pbmig1a)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pbmig1b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgcill1c)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgcill1b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pglaua)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pbmmax1a)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pbdell1b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pbdell1a)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pccoaom1)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgeup2b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgeup2arev)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgdell2b)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pgdell2a)	caggttaggc	atctctgcca	tcatacaaa	aaaa.....	.....	gaaaaaaa	ttccaaagcc	agaaaagaga	tcgtagtta	attagaagat
pileup_49.msf(pccoaom2)	CAGGT-----	TCATC-----	TCATC-----	TCATC-----	TCATC-----	TCATC-----	TCATC-----	TCATC-----	TCATC-----	TCATC-----
Consensus										
pileup_49.msf(pgeup1crev)	aatggccacc	aacgagag-	-----	-----	-----	-----	-----	-----	-----	-----
pileup_49.msf(pgeup1arev)	aatggccacc	aacgagag-	-----	-----	-----	-----	-----	-----	-----	-----
pileup_49.msf(pgeup1brev)	aatggccacc	aacgagag-	-----	-----	-----	-----	-----	-----	-----	-----

```
pileup_49.msf(pgalblc)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgalbia)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgnigia)      aatgaccacc aacggagag~ ~~~~~
pileup_49.msf(pbmaxib)      aatgaccacc aa~~~~~
pileup_49.msf(pgnigib)      aatgaccacc aacggagag~ ~~~~~
pileup_49.msf(pbmigia)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pbmigib)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgcilic)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgcililb)     aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pglaua)       aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pbmaxia)      aatggccac~ ~~~~~~
pileup_49.msf(pbdellb)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pbdellia)     aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pccoaoim1)    aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgeup2b)      aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgeup2arev)   aatggccacc aacggagag~ ~~~~~
pileup_49.msf(pgdel2b)      aatggccgcc aacggagag~ ~~~~~
pileup_49.msf(pgdel2a)      aatggccgcc aacggagag~ ~~~~~
pileup_49.msf(pccoaoim2)    aatggccgcc aacggagag~ ~~~~~
pileup_49.msf(pccoaoim2)    AATG-CG- ~~~~~~
                               Consensus
                               Translation start site
```

**Fig. 14. CCoAOMT promoter-GUS constructs**






Fig. 15. Expression pattern of *pBIN1DC2* (-185) in transgenic poplar.

GUS activity was detected in periderm, cortex and phloem fibres.



Fig. 16. Expression pattern of *pBIN1DA5* in transgenic poplar.

GUS activity was found in cambial ray cells, vessels and ray cells adjacent to the vessels.








Fig. 17. A map of *CCoAOMT* promoters with domains as defined in this invention

+B: positive bark tissue domain; -B: AC-II element, negative bark tissue domain

+CR and -CR: positive and negative cambial ray cell domains

V: vessel and adjacent ray cell domain; CC: companion cell domain

F: phloem fibre domain

W: wounding domain

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Vlaamse Interuniversitair Instituut voor Biotechnologie
- (B) STREET: Rijvisschestraat 118
- (C) CITY: Zwijnaarde
- (E) COUNTRY: Belgium
- (F) POSTAL CODE (ZIP): B-9052
- (G) TELEPHONE: 32 9 244 66 11
- (H) TELEFAX: 32 9 244 66 10

(ii) TITLE OF INVENTION: Tissue-specific poplar promoters

(iii) NUMBER OF SEQUENCES: 2

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (vi) PRIOR APPLICATION DATA

- (A) APPLICATION NUMBER: EP.97.202.507.6
- (B) FILING DATE: 13-AUG-1997

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1994 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Populus trichocarpa

## (ix) FEATURE:

- (A) NAME/KEY: CAAT\_signal
- (B) LOCATION:1841..1848

## (ix) FEATURE:

- (A) NAME/KEY: TATA\_signal
- (B) LOCATION:1871..1876

## (ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION:1..1994

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```
TCTAGAGTTC GTGGTTTATT TTTATATTTT TAAAGTATTT AAAAAAAAAAT AATTTTTTTTA      60
ATTTTTTATT TGGTTTAAAT TAATTTTTTT TTGTATTTT ATATTATTTT GATGTGATGA      120
TATAAAAAAT AAAATTTTAA AAGTAACCCT TGTCTAAATA CAAATAGGC TCTATATATA      180
```

AGCCAGAGGA GGTGATATTT TGATTATTTT CTTTAAGACT ACAGATGACC CGATAACATG	240
AAACAAAAAT ATAAAAACAA GGTCAACCGT GTGACTTGTT CCGCCCCCGT CCCGGGTTCG	300
ACCCTCTATG TGCACGCCTG TCACCCCCGC GGTGCCTTAC CTGCTCCTGG GCTTGCAGGA	360
TGTCCAGTGG GCCGTGGGGA ATAGTCGTGG TGCGCGTAAG CTGGCCCGGA CACCCCATGT	420
AAATCAAAAA AAAAAAAAAA AAAAAACAA GGTACGCGG GCATTCTAG CTTCTCTATA	480
GGCATTTTGC TTTGAGCCTC CGACCATTCC CCCCTGCTCT TTTCTTATTC ACGATAGAAG	540
TCTGTTTCAGT TAAAGAAAAC ATGGCCTGTT TTCCTAGAAA TTTATCTATA AAAAAGGAAA	600
TTCTTGTTAA AATAAAGAAA AAAATTCAAT TGAAATACAC TTCCAGCTT TCGAATGTTT	660
CTATTATATT TTGGAAAAACA TGTGAAAAAT ATTTTATTAA TGTTCACTTG AAAAAATGGT	720
TATGGTTGCT TTTCAAAGTG TTTTTACTT GGAAATGCTT TAAAATAATA TTTTTTTATT	780
TTTTAAAAAT CATTGTGAT ATCAGCGCAT CAAAATGATT TGAAAACATT AAAAAATATT	840
AATTTAAAC AAAAAATAA AATAAAATAA TTTAAATTTT TTAAATACT TTAAACAT	900
AAAAAAAAAA CAGAATAGAA TTTAGACATA AAAAGTGTTA TTCTCGTCGT TTAATAGCGG	960
GAGGTGGTAA GAAACATGCT AAGCATTCAC AGTTTTGGAT ATTGATTATC CATGTGTAC	1020
CCTAAAAAAT GCCTTTTTCC ATTTTCATGAA ATCCTCATTT AATAGCAATT CTTTAAAGGA	1080
GGAGACTAGA GACAGAGGTG GCTGCTGTCA ACCTAGTCGG TGAATTTAAA CTTCAACCGG	1140
CACATATATG CATACCAAGT ATACCCTTTA CATCTGCCCT AATTAAGACT GTAAAACGGA	1200
TTTGGATTTT TGCCGACAAA GGCTAGTTTG TGGAGAAAAA ACAACGAAGA TAAATTTTTA	1260
GATGACAAAG TCAACAATAG TTCGAGAGAT TCTTTAAAGG ACTCATCCGT TGACGGAGGT	1320
GGCCATATGC TACCAACTCT TGGACGTGGA GTCCCTTTGG TAATTCACC TATCCCTCAC	1380
CCAATTTCTA TTAGCAGTTA GCACATGTAA TTTATGATTG GTGAGCCCAG CACAAATCTT	1440
TTCCAGTTAA ACACATATAT TAATTTATGA TTAATTATTT AATTCTCTCC ACTCTTAACA	1500
AATTAATCAT ACATGGCATA ACATTTTAGC TTTTGATCTC GAGAATCTCT ACCTAACCAT	1560
TGACTTCTTT ACTGTTTCAGG AATCTTAGAA AAAAGGAGGA CAAAAAAAAA TATGCCCAAT	1620
AAATTATTTA AGAATTTGAA CCGATATTTG GTGTCATAGA TCCCAAAAAT GACGCCAGCG	1680
ATGCCTAAGG GAAGGAGTAC CACTAGCCCA CAGCACGATA CGATCACCAA CAAGGTGGGT	1740
CCCATATTTG GTGGGCCAAA AACCACATT ATCCTTCGTC CTAACCTACAG GAACCTCACC	1800
AACCCCTCC CGGTTGGTAG CCGGTCCAGC CTCCCCGCTA CTCCAATTCA AACCGGGCTC	1860
TCATTTCCAA TAAATACCAC CCGCCCTTTA CCATTTTCGA TCAGGTTAGG CATCACTACC	1920
ATCATCAACA AAAAAAAAAA AAAAATCCAA GGCCAAGAAA GAGATCGTAG TTTAATTAGA	1980
AGATATACAC AATA	1994

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1363 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Populus trichocarpa*

## (ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION: 1..1363

## (ix) FEATURE:

- (A) NAME/KEY: TATA\_signal
- (B) LOCATION: 1243..1249

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

TCTAGAGAAC ACGGTTTCAA CCGCGTTTCC AAACATGATT TAAATATACA CGGTATATTT      60
TTGCATTTCA ATAGGGTTTT TGAAAAAGAA ATGAATTTTA TTTATTTATT TTGTTTTAAA      120
TTATTTTTTA GTATTTTGTAG ATCGTTTAA CATGTCAATG TCAAAAAAAAA TTAAGAAAAA      180
TATTATTTCA ATTTATTGT AAGAAAAAAA ACCTTAAAAA ACAATTATTA CCAAACATC      240
AAACTGGCTC TGAAAGTATC TCATAGCATA ATGCACTAAC CAATTATTTA AATTTTCCAT      300
CCTGTCATGG AGAAAGATTC CATGGTTGAA GACTGTATGA TAAGGAAAAA TGTCATGAAC      360
TCATGGTATA AGTAATTTC TATCCAATA CAGCAAGCTT GATGTTAGGT TAGGGTTGAT      420
GGTTGTCTTC TTTCATGGAA ATGTTTTGCC ATGCCACAC GAAACAGGCA AGAAAACCAG      480
ACAATATTAG GAATTGTTTC AATGTATTGA TATTAATAAT AAATTTTAAA ATTAATAAAA      540
TATTATTTTA ATATATTTAT AAATAAAAAA TACTTAAAAA AAACATCTAT TACACTTAAA      600
AAAAACATTA ATTAATTACT GCCTAGCTTT ACTAGAAAAT CCACACACTA ACTGGGCGAT      660
TGAAACTCCA GCCATTTTTA TATATTTGTC CTGTGATTAT CATAGACGGT AAAACGAAAT      720
TGGATTTTTT TATTTTGTG GAGAAAAAAA AAAGAAAATA AATATTGTCA GCAGTAAGAC      780
GGAGAGATTC TTAAAAGGAG TCATCCATTG TCAATGCGGT GGCTACGAGC CACCAACTCC      840
CGTGGAGTCA AATTCCTGAG GACACCTCAC CAACCCCTTA CCCACTTTCT ATTAGCAGCA      900
CATGTAGCCA TCCCAACAA CAAAGTGGTG AGCCCACCAC AATTTTCTAC TCTCTACGAT      960
TTAAATCAAT TACACGTGGC ATAAATGTC GAGCCTTTTA TTCAAGAAA CCAAACCTAA      1020
CACCGTGAAC TTAATTTCTT TCGCAAATAT CTAAATAAG GGTACATGAA TTAAATGTAT      1080
AGAAATTGAA TTAGTGTCG AAACCTAAAA TGACCACTGG ACAAACACCG ATAAGTGGGT      1140
CCCCAAATAC CCACGGTGTC CTAAGAACTC ACCAACCCCC ACCCGGTTGG AAGCCGGTCC      1200

```

AACCACCCCA	CTACTCCGGC	TCAAACCGGA	CTCTCATCTC	CAATAAATAC	CACCTGCCCT	1260
TGCCATTTTC	AATCAGGTCA	GACATCCTTA	CCATCGTCGC	CCCCAGAAAA	ACCTTCCAAC	1320
GCCAGGAAAG	AGAGTATAGT	TTTGTTATAA	GATATACAAA	ATA		1363





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/82, 15/29</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/09188</b> <b>(43) International Publication Date:</b> 25 February 1999 (25.02.99)
<b>(21) International Application Number:</b> PCT/EP98/04988 <b>(22) International Filing Date:</b> 27 July 1998 (27.07.98) <b>(30) Priority Data:</b> 97202507.6 13 August 1997 (13.08.97) EP <b>(71) Applicant (for all designated States except US):</b> VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECH- NOLOGIE [BE/BE]; Rijvisschestraat 118, B-9052 Zwij- naarde (BE). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> CHEN, Cuiying [CN/BE]; Meibloemstraat 143, B-9000 Gent (BE). MEYERMANS, Hugo [BE/BE]; Breugelhoevestraat 25, B-2560 Nijlen (BE). VAN MONTAGU, Marc [BE/BE]; De Stassartstraat 120, B-1050 Brussel (BE). BOERJAN, Wout [BE/BE]; Zomerstraat 44 <sup>b</sup> , B-9270 Laarne (BE). <b>(74) Common Representative:</b> VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE; Rijvisschestraat 118, B-9052 Zwijnaarde (BE).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 6 May 1999 (06.05.99)
<b>(54) Title:</b> TISSUE-SPECIFIC POPLAR PROMOTERS  <b>(57) Abstract</b> <p>The invention concerns the isolation and characterization of DNA sequences representing a caffeoyl-CoA-O-methyltransferase (CCoAOMT) promoter having a biological activity in at least one plant or tree vessel and/or in cells adjacent to said vessel or vessels. The cells adjacent to the vessel are xylem ray cells whereas the vessel can be a differentiating vessel.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



# INTERNATIONAL SEARCH REPORT

International Application No.  
PC 98/04988

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C12N15/82 C12N15/29				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N				
Documentation searched other than minimum documentation, to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
P,X	CHEN, C., ET AL.: "Populus trichocarpa CCoAOMT1 gene, exon 1 to exon 5" EMBL ACCESSION NO. AJ223621, 17 July 1998, XP002086171 see the whole document ---	1		
A	GRIMMIG, B., ET AL.: "Structure of the parsley caffeoyl-CoA O-methyltransferase gene, harbouring a novel elicitor responsive cis-acting element" PLANT MOLECULAR BIOLOGY, vol. 33, January 1997, pages 323-341, XP002054325 see the whole document --- <div style="text-align: center;">-/--</div>	1-10		
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>				
<b>* Special categories of cited documents:</b>				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </td> <td style="width: 50%; vertical-align: top;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </td> </tr> </table>			<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>
<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>			
Date of the actual completion of the international search  <div style="text-align: center;">12 March 1999</div>	Date of mailing of the international search report  <div style="text-align: center;">19/03/1999</div>			
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer  <div style="text-align: center;">Maddox, A</div>			

Form PCT/ISA/210 (second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

International Application No.

CT/EP 98/04988

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MENG, H., ET AL.: "Populus tremuloides caffeoyl-CoA 3-O-methyltransferase mRNA" EMBL SEQUENCE DATABASE, REL.44 13-JUN-1995, ACCESSION NO. U27116, XP002054772 see the whole document ---</p>	1-10
A	<p>YE, Z.-H., ET AL.: "Differential expression of two O-methyltransferases in lignin biosynthesis in Zinnia elegans" PLANT PHYSIOLOGY, vol. 108, 1995, pages 459-467, XP002054326 cited in the application see the whole document ---</p>	1-10
A	<p>CAPELLADES, M., ET AL.: "The maize caffeic acid O-methyltransferase gene promoter is active in transgenic tobacco and maize plant tissues" PLANT MOLECULAR BIOLOGY, vol. 31, 1996, pages 307-322, XP002054327 see page 314 - page 315 ---</p>	1-10
A	<p>FEUILLET, C., ET AL.: "Tissue- and cell-specific expression of a cinnamyl alcohol dehydrogenase promoter in transgenic poplar plants" PLANT MOLECULAR BIOLOGY, vol. 27, 1995, pages 651-667, XP002054328 cited in the application see the whole document ---</p>	1-10
A	<p>BEVAN, M., ET AL.: "Tissue- and cell-specific activity of a phenylalanine ammonia-lyase promoter in transgenic plants" EMBO JOURNAL, vol. 8, 1989, pages 1899-1906, XP002054329 cited in the application see the whole document ---</p>	1-10
A	<p>HAWKINS, S., ET AL.: "Cinnamyl alcohol dehydrogenase: identification of new sites of promoter activity in transgenic poplar" PLANT PHYSIOLOGY, vol. 113, February 1997, pages 321-325, XP002054330 cited in the application see the whole document ---</p>	1-10
	-/--	

# INTERNATIONAL SEARCH REPORT

International Application No.

PC 98/04988

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>HAUFFE, K.D., ET AL.: "Combinatorial interactions between positive and negative cis-acting elements control spatial patterns of 4CL-1 expression in transgenic tobacco"</p> <p>THE PLANT JOURNAL, vol. 4, 1993, pages 235-253, XP002054331 cited in the application see the whole document</p>	1-10
A	<p>EP 0 516 958 A (BAYER AG) 9 December 1992 see page 4, line 1 - line 9</p>	1-5

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04988

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0516958	A	09-12-1992	DE 4117747 A	03-12-1992
			CA 2067317 A	01-12-1992
			JP 5199886 A	10-08-1993
			US 5728570 A	17-03-1998
<hr/>				

Form PCT/ISA/210 (patent family annex) (July 1992)